

CONGENITAL ANOMALIES

Volume 50 · Number 4 · December 2010

CONTENTS

Review Article

- 205 Individual variation in organ histogenesis as a causative factor in the developmental origins of health and disease: Unnoticed congenital anomalies?**

Hiroki Otani, Jun Udagawa, Toshihisa Hatta, Yukiko Kagohashi, Ryuju Hashimoto, Akihiro Matsumoto, Fumio Satow, and Masayuki Nimura

Original Articles

- 212 Maternal dietary n-6/n-3 fatty acid ratio affects type 1 diabetes development in the offspring of non-obese diabetic mice**

Yukiko Kagohashi, Norio Abiru, Masakazu Kobayashi, Michio Hashimoto, Osamu Shido, and Hiroki Otani

- 221 Prevalence and distribution of congenital abnormalities in Turkey: Differences between the prenatal and postnatal periods**

Kazim Oztarhan, Ali Gedikbasi, Dogukan Yildirim, Oguz Arslan, Erdal Adal, Sultan Kavuncuoglu, Sibel Ozbek, and Yavuz Ceylan

- 226 Diet with a low n-6/n-3 essential fatty acid ratio when started immediately after the onset of overt diabetes prolongs survival of type 1 diabetes model NOD mice**

Yukiko Kagohashi and Hiroki Otani

Acknowledgement

- 232 Reviewers**

Japanese Teratology Society 50th Annual Meeting, July 8–10, 2010

Program

- A1 Abstracts**

Society News

- J5 Announcement**